The Global Fruit & Veg Newsletter

n° 18 February 2017

Perfect timing for a national fruit and vegetable nutrition policy

Although a diet rich in fresh fruit and vegetables is considered a cornerstone of good health, Canadians are still consuming less than the 7-10 servings per day recommended in Canada's Food Guide. The studies featured this month illustrate how health behaviours in Canada are effected by various interventions and policies. In fact, shopping frequency, awareness of Canada's Food Guide, and school based interventions are all methods we use to assess how overall consumption is impacted by external variables.

In Minaker's "Food shopping is associated with dietary outcomes in Ontario" she finds that food store selection and BMI are correlated, and synthesized that their findings have potential for relevant food intervention programs. Adding to this study, Drapeau examines how school interventions have the potential to directly increase fruit and vegetable consumption among children. Fernandez and Provencher's article examines one of Health Canada's developed initiatives, Eat Well Campaign: Food Skills (EWC) which promotes family meal planning and preparation. It examines the collaboration between partners from the food retail and health sectors, industry and media and how they worked to leverage resources and expertise and extend the reach and effectiveness of the EWC.

To this end, the Canadian Produce Marketing Association (CPMA and the Canadian Public Health Association) are calling on the federal government to establish a Government of Canada policy statement supporting the goal of increasing the fruit and vegetable consumption of Canadians by 20% by 2020. In fact, Canada is the only G7 country without some form of national fruit and vegetable nutrition policy. This goes beyond Canada's Food Guide and into programs like CPMA's Half Your Plate by providing Canadians with specific resources to promote consumption at both the national and local level. The research in this month's newsletter is supporting evidence to move this policy forward, and strengthens the argument of the importance of using intervention and other policy tools to increase consumption across the population.

Ron Lemaire President, Canadian Produce Marketing Association, CANADA

UITS & VEGGIES

PLATE.CA

HALF YOUR



P. Alvarado • 5 a Day Nicaragua H. Bargen • 5 am Tag Germany S. Barnat • Aprifel • France P. Binard • Freshfel Europe • Belgium S. Carballo • 5 a Day Uruguay • MAES L. DiSogra • United Fresh • USA P. Dudley • United Fresh • New Zealand J. Estradas • 5 a Day Bolivia D. Ferreira • 5 ao dia • Portugal G. Fretes • 5 a Day Paraguay C. Gamboa • Network 5 a Day Costa Rica • Ministry of Health ME. Leão Diogenes Melo • F&V Promotion - INCA Brazil A. Gysi • 5 am Tag • Switzerland P. Harycki • 5 a Day Poland • KUPS H. Huss • CO CONCEPT • Luxemburg Z. Huszti • 5 a Day Hungary Y. Iritani • 5 a Day Japan J. Jalkanen • 5 a Day Finland S. Lauxen • 5 am Tag Germany Corporation S. Lewis • Fruits & Veggies Half Your Plate! • Canada C. Macias • F&V Promotion Program of Cuba • Institute of Nutrition and Food Hygiene A. Moises • 5xday • Mexico M. Penny • 5 a Day Peru E. Pivonka • Fruits & Veggies - More Matters • USA G. Rebnes • 5 a Dagen • Norway J. Rey • 5 a Day Spain • Association for the Promotion of consumption of F&V A. Senior • 5 a Day Colombia • Corporación Colombia Internacional M. Tapia • 5 a Day Foundation Venezuela F. Vio • 5 a day Chile Corporation H. von Bargen • 5 am Tag Germany M. Winograd • 5 a Day Association Argentina



Acknowledgement to 250 contributors since 2006

July/August 2013 : E. Pivonka; KA. Grimm and colleagues; E. Howlett and colleagues; S.Burton and colleagues; TO. Erinosho and colleagues (How to increase F&V intake in adults?)

September 2013 : J. Ogden; AR. Khanolkar and colleagues; EL. Melbye; J. Manganello & KC. Smith (Parenting practices and their consequences on children)

October 2013: M. Bruening; V. Cordeiro Barbosa Filho and colleagues ; AM. Santaliestra-Pasías and colleagues, KA Levin (Behavioral risk factors in adolescents) November 2013: M. Bonnefoy ; M. Södergren & SA. McNaughton ; CE. Neville & JV. Woodside; MS. Lee, YT. Lo & ML. Wahlqvist (F&V intake in older adults) December 2013: T. Gibault; W.Willett (How translate F&V beneficial assets into

efficient consumption practice?)

Editions available in:

English:

www.aprifel.com / www.freshfel.org / www.kauppapuutarhaliitto.fi www.unitedfresh.co.nz / www.5amtag.ch / www.halfyourplate.ca French: www.aprifel.com Spanish: www.5aldia.org

THE GLOBAL FRUIT & VEG NEWSLETTER CONTACTS INFO APRIFEL Agency for the Research and Information on Fruits and Vegetables 4 rue de Trévise 75009 Paris – France GLOBAL FRUIT & VEG NEWSLETTER Secretariat : gfvn@interfel.com

erfel.com www.aprifel.com

Food shopping is associated with dietary outcomes in Ontario

Leia M. Minaker

School of Planning, Faculty of Environment, University of Waterloo, and Propel Centre for Population Health Impact, University of Waterloo, CANADA

The food environment and poor diets in Canada

Canadians typically fail to meet recommended daily guidelines for vegetable and fruit consumption, and many Canadians consume poor quality diets. Dietary behaviours are embedded in social, economic, and physical environments. Current diets are a logical response to the existing retail food environment in Canada, which tend to promote the purchase and consumption of less healthy foods and beverages.

Gaps in knowledge about food shopping and diet

The vast majority of dietary energy consumed in Canada is purchased in food stores as opposed to restaurants. Little is known about why consumers choose to shop for food in the places they do, and how buying food at different types of food retailers is associated with dietary and weight outcomes. These gaps in knowledge are important because if we understand why people shop where they do and how food shopping is associated with dietary intake, both retailers and public health practitioners can create effective strategies to improve the diet quality of Canadians - starting in the food store.

Investigating food shopping motivations and associations with diet and weight

We conducted a study using a population-based survey of 4574 residents living in 2596 households from Kitchener, Cambridge, and Waterloo, Ontario to examine: 1) motivations for food store choice; and 2) how food purchasing is associated with dietary and weight outcomes.

Food shopping motivations

First, we asked main household shoppers from the participating households the top three reasons why they choose a specific type of store when they buy food. The most commonly-reported reasons why shoppers chose a specific supermarket were: proximity to home, work or other daily activities, high food quality, and cheaper food prices. For convenience stores, proximity and convenient hours of operation were the most frequently-reported reasons. Shoppers most often reported choosing farmers' markets based on the high quality of foods and because they wanted to "buy local".

Food shopping diet, and weight-related outcomes

Second, we asked how frequently shoppers buy food from the following types of food outlets: supermarkets, supercenters,

convenience stores, specialty stores, farmers' markets, food banks, home delivery, and food co-ops. We also asked them to report their weight, height and waist circumference according to a standard protocol, and how frequently they consumed fruit and vegetables. For a sub-group (n=1362), we collected food records over two days, which captured detailed information on everything participants ate and drank. Most participants (91%) shopped at a supermarket at least once per week, compared to 16% of participants who shopped at supercenters, 10% who shopped at convenience stores, and 7% who shopped at farmers' markets at least once per week. After adjusting for a number of socio-demographic factors (sex, education, car ownership, and household income), we found that people who shop frequently at supermarkets, farmers' markets and food co-ops consumed fruit and vegetables significantly more frequently than those who shopped at these places less frequently. People who frequently shopped at convenience stores, used home delivery services, and used food banks consumed fruit and vegetables significantly less frequently than those who did not use these food outlets as often. People who often shopped at specialty stores and farmers' markets had lower body mass index and smaller waist circumferences than those who did not use these types of food retail as often, whereas shopping at supercenters was associated with marginally higher body mass index and waist circumference.

Farmers' markets, specialty stores, and convenience stores as important settings to promote fruit and vege-table intake

Shopping frequently at farmers' markets and specialty stores was associated with lower weight and more frequent fruit and vegetable intake in the three cities, whereas shopping frequently at convenience stores was associated with less frequent fruit and vegetable intake. Because this study was cross-sectional, we are not able to say that shopping at farmers' markets or specialty stores cause increased fruit and vegetable consumption. Indeed, it may be that people who enjoy consuming fruit and vegetables shop frequently at the farmers' market because of the high quality of available produce. That said, this study builds on past research showing that in-store marketing, food availability, quality and prices can influence food purchasing, which then affects dietary intake. This study suggests that in particular, intervening in convenience stores to promote fruit and vegetable sales may be a promising direction for future interventions to improve fruit and vegetable intake in Canada.





The Effectiveness of A School-Based Nutrition Intervention on Children's Fruit, Vegetable and Dairy Product Intake

Vicky Drapeau

Department of Physical Education, Université Laval, Québec City, QC Canada G1V 0A6 Research Center of the Quebec University Institute of Cardiology and Pulmonology, Québec City, QC CANADA Institute of Nutrition and Functional Food, Université Laval, Québec City, QC CANADA

Even though obesity is recognized as a complex and multifactorial problem, unhealthy eating habits are one important factor contributing to the emergence of childhood obesity and its healthrelated consequences¹. The last Canadian Health Survey revealed that children's eating habits are not optimal and that fruit and vegetables (F&V) and dairy products (DP) represent the two food groups where the largest proportion of children and adolescents do not meet recommendations².

In addition to having a high nutrient density, F&V and DP could play an important role in childhood obesity treatment and prevention. In one previous study, we observed that among 41 food groups, an increase in whole fruit and low-fat milk intake appeared to be the two specific groups consistently associated with better weight control, however, this was observed in adults³. Acute studies have demonstrated that the incorporation of F&V into children's meals can decrease energy intake by lowering energy density⁴. One intervention study also showed a beneficial effect of increasing F&V and DP on children' eating habits compared to a more restrictive intervention that included a reduction in fat⁵. This is line with studies showing that an increase in F&V or DP often results in a concomitant decrease in unhealthy snacks/foods⁶⁻⁸. Thus, a high consumption of these two food groups appear to represent a useful strategy to prevent or treat obesity. At Laval University, our research group has developed a nutrition program called "Team Nutriathlon" where the objective is to increase and diversify the consumption of F&V and DP in children/adolescents.

What is Team Nutriathlon?

Team Nutriathlon is an eight-week, school-based nutrition program where students are encouraged to reach individual and team goals regarding the quantity and variety of F&V and DP consumption (and dairy alternatives) based on Canada's Food Guide9. The individual goals require students to attain a specific quantity of servings per day while the team goals are based on both quantity and variety. During the program, students are invited to report their daily consumption of F&V and DP for each weekday during the eight-week period. This self-monitoring process permits the production of summary reports every two weeks. These reports are then analyzed every two weeks during a "regulation period" where students meet in teams, read the report

also revise their individual and team achievement objectives and identify strategies in order to maintain or increase their F&V and DP consumption. Thus, the program is designed to develop children's autonomy (i.e. decision-making and control over their food choices) towards the gradual adoption and maintenance of healthy eating habits. What is the program efficacy?

and then analyze their results with respect to the team goals. They

In our first study, we evaluated the effectiveness of the program, which included 404 children from grades five and six (intervention n=242, control n=162), and used paper and pencil to record intakes9. At the end of the intervention and even 10 weeks after the program, Nutriathlon participants showed a significant increase in their F&V and DP consumption compared to children in the control group. These positive results prompted us to investigate the efficacy of the program in adolescents. Since school-based programs which incorporate internet use to prevent obesity in youth have been identified as effective strategies to modify eating habits in the school environment¹⁰⁻¹⁴, a Web-based version of the Team Nutriathlon was developed for this group to record intakes. Therefore, the second study assessed the impact of the Web-based Team Nutriathlon among 282 high school students (intervention n=193, control n=89)15. Accordingly, the Web-based version significantly increased the consumption of F&V and DP in this group of adolescents. Thus, Team Nutriathlon appears to be an effective and innovative approach to promote F&V and DP intake in children and adolescents, at least in the short-term.

What are the next Steps?

Because our previous results have shown that parental involvement is a key factor in the program's success, the next step of this initiative is to evaluate its effectiveness among families that have at least one obese child. Results from our pilot project, which includes 13 families randomized to either Nutriathlon or a control group, has shown that the intervention increased the consumption of F&V and DP in children and the global quality of their diet compared to the control group (Drapeau et al., unpublished). Overall, these results suggest that this program represents a promising, positive and non-restrictive obesity prevention program in school settings as well as a clinical tool used in childhood obesity.

References

1. Must A, Strauss RS. Risks and consequences of childhood and adolescent obesity. Int J Obes Relat Metab Disord 1999;23 Suppl 2:52-11.	Children's Fruit, Vegetables, and Dairy Product Intake. J Sch Health 2016;86(5):353-62.
 Garriguet D. Vue d'ensemble des habitudes alimentaires des canadiens, 2004. In: canadiennes RdlEslsdlc, ed.: Statistique Canada, 2006. Drapeau V et al Modifications in food-group consumption are related to long. 	10. Hamel LM, Robbins LB. Computer- and web-based interventions to promote healthy eating among children and adolescents: a systematic review. J Adv Nurs 2013;69(1):16-30.
term body-weight changes. Am J Clin Nutr 2004;80(1):29-37.	 Di Noia J, et al.Computer-mediated intervention tailored on transtheoretical model stages and processes of change increases fruit and vegetable consumption among urban African-American adolescents. Am J Health Promot 2008;22(5):336- 41. Mauriello LM, et al. Results of a multi-media multiple behavior obesity prevention program for adolescents. Prev Med 2010;51(6):451-6. Winett R, et al The Effects of the Eat4Life Internet-Based Health Behavior
4. Pourshahidi LK, et al. Livingstone MB. Influencing and modifying children's energy intake: the role of portion size and energy density. Proc Nutr Soc 2014;73(3):307-406	
5. Epstein LH, et al. Increasing fruit and vegetable intake and decreasing fat and sugar intake in families at risk for childhood obesity. Obes Res 2001;9(3):171-8.	
6. Iak NI, et al. Ine effects of a fruit and vegetable promotion intervention on unhealthy snacks during mid-morning school breaks: results of the Dutch Schoolgruiten Project 1 Hum Nutr Diet 2010;33(6):609-15	Program on the Nutrition and Activity Practices of High School Girls. Journal of Gender, Culture and Health 1999;4(3):239-54.
7.Bere E, et al. One year of free school fruit in Norway7 years of follow-up. Int J Behav Nutr Phys Act 2015;12:139.	14. Randi Schoenfeld E, et al. Using the internet to educate adolescents about osteoporosis: application of a tailored web-education system. Health Promot Pract 2010;11(1):104-11.
8. Andersen LB, et al. The effects of water and dairy drinks on dietary patterns in overweight adolescents. Int J Food Sci Nutr 2016;67(3):314-24.	15. Chamberland K, et al. The impacts of a computer-assisted school-based nutrition intervention on the consumption of vegetables, fruits, and dairy products in

9. Drapeau V. et al. The Effectiveness of A School-Based Nutrition Intervention on

intervention on the consumption of vegetables, fruits, and dairy products in adolescents, ISBNPA online abstract book 2015:190:335.



Selecting cross-sector partners to work with the federal government on promoting healthy eating

Melissa Anne Fernandez and Véronique Provencher

Institute of Nutrition and Functional Foods, Université Laval, Pavillon des Services, Quebec (Quebec) G1C 0A6 CANADA School of Nutrition, Université Laval, Pavillon Paul-Comtois, Quebec (Quebec), CANADA

In response to child overweight and obesity being made a public health priority, Health Canada developed a series of healthy eating and education awareness initiatives¹. The final and third initiative was a social marketing campaign, the Eat Well Campaign: Food Skills (EWC), that promoted family meal planning and preparation to Canadian parents. Health Canada collaborated with cross-sector partners from the food retail industry, the media and the health sector to extend the reach and effectiveness of the EWC. Leveraging resources and expertise are important incentives for governments to work with partners. Developing solutions to address complex health problems, such as improving dietary behaviors, is complicated and requires the joint action of multiple sectors from the government, private industry and civil society^{2,3}.

The adoption of healthy eating initiatives by cross-sector partners, and motivations for organizations from different sectors to collaborate with a federal body have not been examined. Given that adoption can influence an intervention's effectiveness⁴, it is important to understand the factors involved, particularly in the context of adopters from multiple sectors. The objective of this qualitative study was to describe factors that influenced the adoption of the EWC. Findings were based on telephone interviews with 18 of Health Canada's partners. Five main recommendations for public health organizations were derived from the analysis of these interviews⁵.

1. Use a targeted approach to recruit partners

The adoption rate among cross-sector partners appeared to be extremely high. All partners that were directly invited to participate in the EWC agreed to partner with Health Canada. These partners were intentionally sought out by Health Canada through specific networks or invitations to respond to calls for proposals. Pre-selecting and targeting partners that will likely respond well to an initiative's objectives appears to be a very effective method for recruiting partners.

2. Find partners that are highly compatible

All partners that agreed to adopt the EWC had some level of compatibility with the campaign's initiatives, values, audience or staff. Compatibility is a known factor that drives adoption of an initiative by an organization⁶. Having a good fit, particularly when it comes to organizational values, appears to be more important than profit oriented or non-health related missions such as entertainment. We recommend identifying partners that would have a good innovation-systems fit with the public health initiative⁷.

3. Use social networks (peer pressure) with high opinion leadership

Using established networks to recruit organizations provided an automatic partnership base. Some organizations participated, because it was part of their mandate within a specific network, while others felt it would have been impossible to say no given that their entire professional network was expected to adopt the EWC. Using networks with high opinion leadership value is recommended to recruit partners⁸.

4. Use the agency's reputation as leverage

Collaborating with or being associated with Health Canada was a major perceived relative advantage of the EWC. Organizations felt that working with a reputable organization would help improve their credibility and social image among their audience or clients. Relative advantages are among the most important factors influencing adoption. Social prestige, of working with Health Canada, is a prominent sub-dimension of relative advantages that is known to have a strong impact on adoption⁹. Prominent agencies with high levels of credibility can easily leverage their reputation to recruit the most desirable partners.

5. Understand adoption barriers

Only partners that were invited through a third party appeared to reject the EWC. Nothing, however, is know about this group of nonadopters, because they were not made known by the third party to Health Canada and could not be included in the study. Health Canada speculated that non-adoption was likely due to organizations having a smaller size with fewer resources and less capacity to collaborate on initiatives outside of their day-to-day business. Not having any information about this group's reasons for rejecting the EWC has left a knowledge gap unfilled about non-adoption, thus it recommended to routinely collect information about reasons for not adopting an initiative. This is particularly important to better understand how to include a diversity of organizations to ensure multiple sectors implicate themselves and are represented in concerted public health initiatives.

Conclusions

Taking into consideration the different needs and expectations of organizations from diverse sectors is important when collaborating with cross-sector partners. Despite the probable differences of organizational missions between partners from the food retail industry, the media and the health sector, there were commonalities among these partners that led to collaborating with Health Canada and adopting the EWC such as shared values. In Canada, prominent federal bodies like Health Canada with strong reputations are desirable partners and these organizations can leverage their notoriety to attract the highest quality partners to help advance population health goals.

References

- 1. Public Health Agency of Canada. Curbing childhood obesity: An overview of the federal, provincial and territorial framework for action to promote healthy weights. Ottawa: PHAC; 2010.
- 2. Riley BL et al. Knowledge to action for solving complex problems: Insights from a review of nine international cases. Health promotion and chronic disease prevention in Canada. 2015;35(3):47-53.

 Vandenbroek P, Goossens J, Clemens M. Tackling obesities: Future choices - buliding the obesity systems map UK: Government Office for Science; 2007 [cited 2015 July 27]. Available from: https://www.gov.uk/government/publications/reducing-obesity-obesity-system-map.
 Virginia Tech. Re-aim: A model for the planning, evaluation, reporting and review of translational research and practice Blacksburg, VA: Virginia Polytechnic Institute and State University; 2014 [cited 2015 July 29]. Available from: http://www.re-aim.hnfe.vt.edu/. 5. Fernandez M et al. Factors influencing the adoption of a healthy eating campaign by federal cross-sector partners: A qualitative study. BMC public health. 2016;In Press.

 Tornatzky LG, Klein KJ. Innovation characteristics and innovation adoption-implementation: A meta-analysis of findings. IEEE Transactions on Engineering Management. 1982;EM-29(1):28-45.

7. Greenhalgh T et al. Diffusion of innovations in service organizations: Systematic review and recommendations. Milbank Q. 2004;82(4):581-629.

8. Dearing JW. Improving the state of health programming by using diffusion theory. J Health Commun. 2004;9 Suppl 1:21-36.

9. Greenhalgh T. Diffusion of innovations in health service organisations a systematic literature review. 2005.

